CUMULATIVE AUTHOR INDEX
Annals of the Entomological Society of America
Volume 100, 2007

A

Abbot, P., 887
Abrahamson, W. G., 539
Aguin-Pombo, D., 19
Ahmad, I., 673, 830
Alarie, Y., 655, 787
Askew, R. R., 381
Azizi, T., 270

B

Backus, E. A., 296
Barbour, J. D., 333
Baselga, A., 444
Bayram, A., 574
Bell, A. J., 876
Bidau, C. J., 850
Billick, I., 887
Birge, L. M., 721
Bouchard, P., 178
Braswell, W. E., 721
Brewer, M. J., 196
Brodie, E. L., 345
Buck, M., 9
Burns, S. N., 257
Bush, G. L., 474

C

Caceres, C., 900
Calcaterra, L. A., 470, 512
Calvert, F., 433
Cane, J. H., 350
Cassill, D. L., 327
Cave, R. D., 164, 839
Chan, M.-L., 359
Chang, C. L., 562, 900
Chee-Sanford, J., 275
Chen, G., 245
Chen, J.-X., 36
Chen, X.-S., 366
Chen, X.-X., 390, 425
Chenoweth, L. B., 311
Choe, D.-H., 320
Clarkin, C. E., 539
Cline, A. R., 296
Cognato, A. I., 582
Connor, E. F., 375
Coombs, M. T., 736
Cooper, W. R., 236
Cope, N., 339
Corak, I., 574
Costa, H. S., 736
Coulson, R. N., 289
Courtney, G. W., 91
Cranston, P. S., 617
Cronin, J. T., 640
Crossno, J. E., 474
Cueva del Castillo, R., 591

D

Danisman, T., 574
Davis, A. K., 339
De Barro, P., 736
de Guzman, L. I., 187
de León, J. H., 749
de Melo Sene, F., 490
de Remes Lenicov, A. M. M., 798
Dillwith, J. W., 83
Dong, P., 228
Dorchin, N., 539
Dosdall, L. M., 178, 861
Downer, K. E., 251
Du, J., 245
Duncan, R. E., 413

E

Edde, P. A., 83
Eitam, A., 603
Ekesi, S., 562, 900
Ellerseick, M. R., 296
Espírito-Santo, M. M., 95

F

Fasnacht, M. P., 893
Feng, M.-G., 425
Fernandes, G. W., 95
Ferris, V. R., 649
Florane, C., 418
Follett, P. A., 433
Forschler, B. T., 763
Foster, P. G., 27
Frake, A. M., 187
Franquinho Aguiar, A. M., 19
French, B. W., 59

G

Gaimari, S. D., 196
Galindo-Cardona, A., 711
Gavrilovic, V., 474
Gibson, G.A.P., 861
Gilbert, G., 609
Giray, T., 711
Golden, M., 433
Gómez, J. F., 381
Gray, D. R., 439
Griswold, T., 350

H

Hajek, A. E., 876
Hall, H. G., 64
Hall, J. P. W., 773
Hammack, L., 59
Hammer, S., 887
Hancock, E. G., 914
Hanks, L. M., 333
Hannam, J. J., 876
Harris, M. K., 582
Hartsough, C. D. B., 375
Hass, M., 53
Hasson, E., 490
Hauser, M., 152
Haverty, M. I., 206
Hazel, W., 53
He, J.-H., 390
Heath, M. S., 483
Henderson, G., 506, 703
Hijii, N., 222, 869
Horton, D. R., 403
Howard, D. J., 721
Hu, F., 228
Huang, Z., 245

I

Ikai, N., 869
Ingber, B., 418
Inouye, B. D., 596
Isaias, R. M. D. S., 907

J

Jacas, J.-A., 413
James, D. G., 41
Japoshvili, G., 683
Jayasimha, P., 506, 703
Jepsen, S. J., 549
Jia, F.-X., 228
Johnston, J. S., 270, 289
Judd, T. M., 893

K

Kaiser, M. E., 196
Kandemir, I., 289
Keddie, B. A., 861
Kim, J.-P., 1
Klimov, P. B., 810
Koehler, P. G., 568
Korkmaz, M., 265
Kubota, K., 100
Kuznetsova, V. G., 19

L

Lacey, E. S., 333
Lamas, G., 449
Lechner, M. G., 345
Lee, C.-F., 623
Lee, M.-L., 64
Legaspi, B. C., Jr., 497

CUMULATIVE AUTHOR INDEX
Annals of the Entomological Society of America
Volume 100, 2007

A

Abbot, P., 887
Abrahamson, W. G., 539
Aguin-Pombo, D., 19
Ahmad, I., 673, 830
Alarie, Y., 655, 787
Askew, R. R., 381
Azizi, T., 270

B

Backus, E. A., 296
Barbour, J. D., 333
Baselga, A., 444
Bayram, A., 574
Bell, A. J., 876
Bidau, C. J., 850
Billick, I., 887
Birge, L. M., 721
Bouchard, P., 178
Braswell, W. E., 721
Brewer, M. J., 196
Brodie, E. L., 345
Buck, M., 9
Burns, S. N., 257
Bush, G. L., 474

C

Caceres, C., 900
Calcaterra, L. A., 470, 512
Calvert, F., 433
Cane, J. H., 350
Cassill, D. L., 327
Cave, R. D., 164, 839
Chan, M.-L., 359
Chang, C. L., 562, 900
Chee-Sanford, J., 275
Chen, G., 245
Chen, J.-X., 36
Chen, X.-S., 366
Chen, X.-X., 390, 425
Chenoweth, L. B., 311
Choe, D.-H., 320
Clarkin, C. E., 539
Cline, A. R., 296
Cognato, A. I., 582
Connor, E. F., 375
Coombs, M. T., 736
Cooper, W. R., 236
Cope, N., 339
Corak, I., 574
Costa, H. S., 736
Coulson, R. N., 289
Courtney, G. W., 91
Cranston, P. S., 617
Cronin, J. T., 640
Crossno, J. E., 474
Cueva del Castillo, R., 591

D

Danisman, T., 574
Davis, A. K., 339
De Barro, P., 736
de Guzman, L. I., 187
de León, J. H., 749
de Melo Sene, F., 490
de Remes Lenicov, A. M. M., 798
Dillwith, J. W., 83
Dong, P., 228
Dorchin, N., 539
Dosdall, L. M., 178, 861
Downer, K. E., 251
Du, J., 245
Duncan, R. E., 413

E

Edde, P. A., 83
Eitam, A., 603
Ekesi, S., 562, 900
Ellerseick, M. R., 296
Espírito-Santo, M. M., 95

F

Fasnacht, M. P., 893
Feng, M.-G., 425
Fernandes, G. W., 95
Ferris, V. R., 649
Florane, C., 418
Follett, P. A., 433
Forschler, B. T., 763
Foster, P. G., 27
Frake, A. M., 187
Franquinho Aguiar, A. M., 19
French, B. W., 59

G

Gaimari, S. D., 196
Galindo-Cardona, A., 711
Gavrilovic, V., 474
Gibson, G.A.P., 861
Gilbert, G., 609
Giray, T., 711
Golden, M., 433
Gómez, J. F., 381
Gray, D. R., 439
Griswold, T., 350

H

Hajek, A. E., 876
Hall, H. G., 64
Hall, J. P. W., 773
Hammack, L., 59
Hammer, S., 887
Hancock, E. G., 914
Hanks, L. M., 333
Hannam, J. J., 876
Harris, M. K., 582
Hartsough, C. D. B., 375
Hass, M., 53
Hasson, E., 490
Hauser, M., 152
Haverty, M. I., 206
Hazel, W., 53
He, J.-H., 390
Heath, M. S., 483
Henderson, G., 506, 703
Hijii, N., 222, 869
Horton, D. R., 403
Howard, D. J., 721
Hu, F., 228
Huang, Z., 245

I

Ikai, N., 869
Ingber, B., 418
Inouye, B. D., 596
Isaias, R. M. D. S., 907

J

Jacas, J.-A., 413
James, D. G., 41
Japoshvili, G., 683
Jayasimha, P., 506, 703
Jepsen, S. J., 549
Jia, F.-X., 228
Johnston, J. S., 270, 289
Judd, T. M., 893

K

Kaiser, M. E., 196
Kandemir, I., 289
Keddie, B. A., 861
Kim, J.-P., 1
Klimov, P. B., 810
Koehler, P. G., 568
Korkmaz, M., 265
Kubota, K., 100
Kuznetsova, V. G., 19

L

Lacey, E. S., 333
Lamas, G., 449
Lechner, M. G., 345
Lee, C.-F., 623
Lee, M.-L., 64
Legaspi, B. C., Jr., 497

Legaspi, J. C., 497
Lehman, R. M., 275
Levine, E., 522
Lewis, T. M., 403
Li, C., 27
Li, S., 649
Li, Z.-Z., 366
Liebherr, J. K., 876
Liu, S., 425
Livore, J. P., 512
Logan, J. A., 439
Lopez, J., 394
Lugo-Olguín, S. D., 591
Lundgren, J. G., 275
Luongo, M. P., 539

M
Machado, V., 394
Manfrin, M. H., 490
Marcos-García, M. A., 914
Mariani, R., 798
Marshall, S. A., 9
Martí, D. A., 850
Matsuura, K., 532
Matthews, C. E., 549
McPherson, J. E., 673
Meagher, R. L., 394
Mellencamp, K., 53
Michat, M. C., 655
Miller, K. B., 787
Montllor Curley, C., 345
Moreira, G. R. P., 907
Morel, L., 758
Morgan, D. J. W., 749
Munson, A. S., 439
Mureb Sallum, M. A., 27
Murphy, C. 418
Murphy, N., 1
Mynhardt, G., 582

N
Nachman, R. J., 251
Nagoshi, R. N., 394
Nderitu, P. W., 562
Nelson, L. J., 206
Nielsen, C., 876
Nieves-Aldrey, J. L., 381
Noma, T., 196
Novoa, F., 444

O
Oconnor, B. M., 810
Oku, K., 69
Olfert, O., 861
Oseto, C. Y., 649

P
Palacios-Vargas, J. G., 36
Panizzi, A. R., 745
Park, Y. C., 1
Park, Y. I., 418
Parker, F. D., 350
Patton, J. C., 289
Peña, J. E., 413
Pensel, S. M., 283
Pescador, M. L., 173
Petanovic, R. U., 157
Peters, J. G., 173
Phillips, T. W., 83
Pike, K. S., 196
Pilgrim, E. M., 688
Pinto, M. A., 289
Pitts, J. P., 512
Powell, J. A., 92
Prischmann, D. A., 41
Purcell, A. H., 345

R
Raina, A., 418
Rector, B. G., 157
Reithel, J. S., 887
Remis, M. I., 283
Ren, D., 666
Reyes-Castillo, P., 711
Ricarte, A., 914
Rieske, L. K., 236
Rinderer, T. E., 187
Robertson, J. B., 83
Rodrigues, D., 907
Roehrdanz, R. L., 522
Rosell, R. C., 736
Rosenheim, J. A., 549
Rosenheim, M. R., 549
Rotheray, G. E., 914
Rubink, W. L., 289
Rust, M. K., 320

S
Sabat, A. M., 711
Sak, O., 265
Sampaio, D. S., 907
Sanborn, A. F., 483
Sasakawa, K., 100
Schaefer, C. W., 830
Schiff, N. M., 289
Schwarz, D., 474
Schwarz, M. P., 1, 311
Scott, E. R., 539
Şengül, Ş. H., 265
Serrano, M. S., 296
Sheppard, W. S., 289
Shi, P., 245
Silva, F. A. C., 745
Silvie, P., 394
Sites, R. W., 139
Sithiprasasna, R., 27
Smith, A., 339
Smith, D. R., 375
Smith, D. S., 549
Smith, J. A., 568
Smith, J. J., 474
Smith, T. R., 164, 839
Snyder, W. E., 41
Solensky, M. J., 339
Sosa, A. J., 798
Soto, I. M., 490
Spicer, G. S., 375
Stark, R., 53
Stoffolano, J. G., Jr., 251
Storm, C. P., 41
Sun, L., 649

T
Takakura, K. I., 729
Tan, J., 666
Tanner, J. A., 439
Tauber, C. A., 110
Teal, P.E.A., 257
Thiele, R., 596
Thomsen-Archer, K., 403
Tsutsui, N. D., 512

U
Uçkan, F., 265
Ulmer, B. J., 178, 861
Unruh, T. R., 403

V
Vander Meer, R. K., 257, 512, 758
Vargas, R. I., 603
Vinson, S. B., 270, 327
Virant-Doberlet, M., 73
Vitheepradit, A., 139
Vockeroth, J. R., 196
von Dohlen, C. D., 688

W
Wan, Z.-W., 425
Wang, J.-J., 228
Wang, Y.-P., 390
Werne, A., 53
Wheeler, T. A., 609
Whiting, M. F., 787
Whitman, J. G., 763
Wilkerson, R. C., 27
Williams, K., 418
Willmott, K. R., 449
Wilson, S. W., 640
Winterton, S. L., 152
Wright, L. C., 41

X
Xu, P., 425

Y
Yamashita, S., 222
Yamazaki, K., 729
Yang, J.-T., 359
Yang, L., 366
Yano, S., 69
Yashiro, T., 532
Yigit, N., 574
Yoo, J.-S., 1

Z
Žežlina, I., 73
Zhang, F., 36

CUMULATIVE KEY WORD INDEX
Annals of the Entomological Society of America
Volume 100, 2007

1

18S rDNA,
phylogenetic analysis, 649
restriction fragment-length polymorphism, 649
sunflower seed weevil, 649

5

5-Aza-dC,
Achroia grisella, 265
Apanteles galleriae, 265
risk assessment, 265
toxicity, 265

A

Achroia grisella,
5-Aza-dC, 265
Apanteles galleriae, 265
risk assessment, 265
toxicity, 265
acoustic,
biology, 483
key for identification, 483
new species, 483
taxonomy, 483
Acroceridae,
Baltic amber, 152
Diptera, 152
fossil, 152
Nemestrinoidea, 152
adaptation,
artificial diet, 562
Bactrocera invadens, 562
rearing, 562
Africanized honey bee,
Apis mellifera, 289
cytochrome *b*, 289
mitochondrial DNA, 289
mitotype, 289
aggregation pheromone,
beetle size, 83
host plant effect, 83
pheromone production, 83
plant species, 83
allogrooming,
feeding, 763
social interactions, 763
subterranean termite, 763
worker behavior, 763
Allonemobius,
hybridization, 721
Orthoptera, 721
sympatry, 721
temporal partitioning, 721
Alydidae,
Caribbean, 830
Hemiptera, 830
Heteroptera, 830
Neotropical, 830
Anagrus,
blackberry, 41
Erythroneura, 41
grape, 41
leafhoppers, 41
Anagrus nigriventris,
egg parasitoid, 640
planthopper, 640
prairie potholes, 640
Scolochloa festucacea, 640
Andes,
Ithomiinae, 449
taxonomy, 449
Anisoscelini,
insect–plant interactions, 907
Passifloraceae, 907
scanning electron microscopy, 907
stylet penetration, 907
Anopheles,
COI, 27
Leucosphyrus Group, 27
ND6, 27
phylogeny, 27
ant,
micronutrients, 893
Myrmica, 893
Reticulitermes, 893
termite, 893
ant–aphid interactions,
mutualisms, 887
predation, 887
Anza Borrego Desert State Park,
brittlebush, 320
feeding deterrent, 320
Forelius pruinosus, 320
Linepithema humile, 320
Apanteles galleriae,
5-Aza-dC, 265
Achroia grisella, 265
risk assessment, 265
toxicity, 265
Aphis glycines,
biological control, 196
natural enemies, 196
Apidae,
biogeography, 810
Centriacarus, 810
Chaetodactylidae, 810
Megachilidae, 810
Apinae,
bees, 596
Centridini, 596
oligolecty, 596
pollination, 596
Apis mellifera, 187, 289
Africanized honey bee, 289
cytochrome *b*, 289
instantaneous growth rate, 187
mitochondrial DNA, 289
mitotype, 289
resistance, 187
Russian honey bees, 187
Varroa destructor, 187
Apoidea,
bees, 350
nest, 350
trap-nest, 350
Araneae,
Arctosa, 1
Lycosidae, 1
molecular phylogeny, 1
wolf spider, 1
Arctosa,
Araneae, 1
Lycosidae, 1
molecular phylogeny, 1
wolf spider, 1
artificial diet,
adaptation, 562
Bactrocera invadens, 562
rearing, 562
Aspidobracon,
Braconidae, 390
China, 390
new species, 390
Philomacroploea, 390
Auchenorrhyncha,
chorus, 73
mating behavior, 73
nocturnal activity, 73
vibrational communication, 73
avian predation,
edge effect, 729
fleeing behavior, 729
habitat structure, 729
urbanization, 729
azadirachtin,
hemolymph, 245
lipid metabolism, 245
Ostrinia furnacalis, 245
protein expression, 245

B

bacteria,
mating, 418
oviposition, 418
primary reproductives, 418
ultrastructure, 418
bacteriocytes,
Bemisia aff. *gigantea*, 736
egg maturation, 736
endosymbionts, 736
pharate adults, 736
Bactrocera invadens,
adaptation, 562
artificial diet, 562
rearing, 562
Baltic amber,
Acroceridae, 152
Diptera, 152
fossil, 152
Nemestrinoidea, 152
basitarsal pad,
bean leaf beetle, 59
corn rootworm beetle, 59
sexual dimorphism, 59
Baumannia,
endosymbionts, 345
glassy-winged sharpshooter, 345
Homalodisca, 345
Wolbachia, 345
bean leaf beetle,
basitarsal pad, 59
corn rootworm beetle, 59
sexual dimorphism, 59
bee nest,
Cerioidini, 914
immature stages, 914
sap run, 914
tree hole, 914
bees, 350, 596
Apinae, 596
Apoidea, 350
Centridini, 596
nest, 350
oligolecty, 596
pollination, 596
trap-nest, 350
beetle size,
aggregation pheromone, 83
host plant effect, 83
pheromone production, 83
plant species, 83
beetles,
Cybocephalus, 839
Pycnocephalus, 839
scale predator, 839
taxonomy, 839
behavior,
development, 425
immature stage, 425
Opius caricivorae, 425
Bemisia aff. *gigantea*,
bacteriocytes, 736
egg maturation, 736
endosymbionts, 736
pharate adults, 736
Berchmansus,
Gonzaga, 110
Leucochrysa, 110
new tribal assignments, 110
Vieira, 110
Bergmann's rule,
clinal variation, 850
Dichroplus pratensis, 850
Rensch's rule, 850
sexual size dimorphism, 850
biocontrol,
egg–pupal parasitoid, 603
host location, 603
Tephritidae, 603
biogeography, 206, 810
Apidae, 810
Centriacarus, 810
Chaetodactylidae, 810
chemotaxonomy, 206
Megachilidae, 810
Reticulitermes tumiceps, 206
speciation, 206
subterranean termite, 206
biological control, 196, 275, 413, 470, 506, 512, 749, 876
Aphis glycines, 196
Coptotermes formosanus, 506
decapitating fly, 470
diagnostic markers, 749
Diaprepes abbreviatus, 413
dispersal, 506
egg parasitoid, 413
endosymbiont, 275
field crop, 876
fire ant, 470
fire ants, 512
geographical distribution, 512
glassy-winged sharpshooter, 749
Glocophyllum trabeum, 506
Gonatocerus walkerjonesi, 749
granivore, 275
insect, 275
invasive species, 876
natural enemies, 196, 749
parasitoid, 470
pitfall trapping, 876
predator communities, 876
Pseudacteon obtusus, 512
Solenopsis gayi, 512
South America, 470
suppression, 506
biological control of weeds,
Dipsacus, 157
Eriophyidae, 157
Leipothrix, 157
systematics, 157
biology,
acoustic, 483
key for identification, 483
new species, 483
taxonomy, 483
blackberry,
Anagrus, 41
Erythroneura, 41
grape, 41
leafhoppers, 41
body size-related traits,
Orthoptera, 283
pericentric inversion, 283
sexual selection, 283
Braconidae,
Aspidobracon, 390
China, 390
new species, 390
Philomacroploea, 390
brittlebush,
Anza Borrego Desert State Park, 320
feeding deterrent, 320
Forelius pruinosus, 320
Linepithema humile, 320
Brochymena,
Halyini, 673
male genitalia, 673
Parabrochymena, 673

C

cactophilic *Drosophila*,
developmental time, 490
host plants, 490
phenotypic plasticity, 490
viability, 490
cactus moth,
development, 497
fecundity, 497
life history, 497
Opuntia cacti, 497
Canary Islands,
exotic species, 19
leafhoppers, 19
Madeira, 19
Sophonia orientalis, 19
canopy,
Cecidomyiidae, 95
global richness, 95
insect galls, 95
tropical rain forest, 95
Caribbean,
Alydidae, 830
Hemiptera, 830
Heteroptera, 830
Neotropical, 830
Carnidae,
new genus, 9
phylogeny, 9
protandrium, 9
symmetry, 9

Carya,
hickory, 582
molecular systematics, 582
North America, 582
pest species, 582
Catiniidae,
Jiulongshan Formation, 666
new genus, 666
new species, 666
Yixian Formation, 666
Cecidomyiidae,
canopy, 95
global richness, 95
insect galls, 95
tropical rain forest, 95
Centriacarus,
Apidae, 810
biogeography, 810
Chaetodactylidae, 810
Megachilidae, 810
Centridini,
Apinae, 596
bees, 596
oligolecty, 596
pollination, 596
Ceratitis capitata,
hydrolyzed yeast, 900
liquid diet, 900
whole cell yeast, 900
Cerioidini,
bee nest, 914
immature stages, 914
sap run, 914
tree hole, 914
Ceutorhynchus neglectus,
Ceutorhynchus obstrictus, 861
courtship behavior, 861
parasitoid development, 861
spatio-temporal distribution, 861
Ceutorhynchus obstrictus,
Ceutorhynchus neglectus, 861
courtship behavior, 861
parasitoid development, 861
spatio-temporal distribution, 861
Ceutorhynchus subpubescens,
Descurainia sophia, 178
flixweed, 178
larval taxonomy, 178
stem mining, 178
Chaetodactylidae,
Apidae, 810
biogeography, 810
Centriacarus, 810
Megachilidae, 810
Chalcidoidea,
Idiomacromerus, 381
Iraella, 381
opium poppy, 381
Torymidae, 381
chemoreception,
contact pheromone, 333
mating behavior, 333
Prionus californicus, 333
chemotaxonomy,
biogeography, 206
Reticulitermes tumiceps, 206
speciation, 206
subterranean termite, 206
chestnut,
Curculio, 236
endophyte, 236
Ormyrus, 236
Torymus, 236
China,
Aspidobracon, 390
Braconidae, 390
new species, 390
Philomacroploea, 390
cholecystokinin,
engorgement, 251
neuropeptides, 251
satiety, 251
sulfakinins, 251
chorus,
Auchenorrhyncha, 73
mating behavior, 73
nocturnal activity, 73
vibrational communication, 73
Cicadellidae,
Mohunia, 366
Mukariinae, 366
new genera, 366
new species, 366
clinal variation,
Bergmann's rule, 850
Dichroplus pratensis, 850
Rensch's rule, 850
sexual size dimorphism, 850
COI,
Anopheles, 27
Leucosphyrus Group, 27
ND6, 27
phylogeny, 27
Coleoptera,
Dytiscidae, 655
Laccornellus, 655
larvae, 655
phylogeny, 655
Coleotroctellus,
new record, 359
new species, 359
Taiwan, 359
Troctopsocidae, 359
Collembola,
Entomobryidae, 36
Mexico, 36
new species, 36
Willowsia, 36
colonization,
island ecology, 711
limited mobility, 711
color polymorphism,
inheritance, 433
southern green stink bug, 433
X-linked trait, 433
competition,
mushroom growth pattern, 222
mycophagous insect community, 222
Nitidulidae, 222
preference-performance linkage, 222
conditional strategy,
pupal color, 53
pupation, 53
stemmata, 53
contact pheromone,
chemoreception, 333
mating behavior, 333
Prionus californicus, 333
Coptotermes formosanus, 506, 703
biological control, 506
dispersal, 506
dual culture techniques, 703
Gloeophyllum trabeum, 506, 703
gut cultures, 703
integument cultures, 703
suppression, 506
corn rootworm,
*fts*Z, 522
mtDNA, 522
Wolbachia, 522
wsp, 522
corn rootworm beetle,
basitarsal pad, 59
bean leaf beetle, 59
sexual dimorphism, 59
cotton,
egg deposition, 745
insect rearing, 745
soybean, 745
stink bug, 745
courtship behavior,
Ceutorhynchus neglectus, 861
Ceutorhynchus obstrictus, 861
parasitoid development, 861
spatio-temporal distribution, 861
cryptic species,
genitalia, 403
geographic variation, 403
morphometrics, 403
mtDNA, 403
Curculio,
chestnut, 236
endophyte, 236
Ormyrus, 236
Torymus, 236
Cybocephalus, 164, 839
beetles, 839
Pycnocephalus, 164, 839
revision, 164

scale predator, 839
taxonomy, 164, 839
cysteine protease inhibitor,
flight muscle, 270
red imported fire ant queens, 270
cytochrome *b*,
Africanized honey bee, 289
Apis mellifera, 289
mitochondrial DNA, 289
mitotype, 289

D

Danaus plexippus,
mating success, 339
wing color, 339
dealation,
juvenile hormone, 257
nuptial flight, 257
precocene, 257
decapitating fly,
biological control, 470
fire ant, 470
parasitoid, 470
South America, 470
Descurainia sophia,
Ceutorhynchus subpubescens, 178
flixweed, 178
larval taxonomy, 178
stem mining, 178
development, 228, 425, 497
behavior, 425
cactus moth, 497
fecundity, 497
immature stage, 425
life history, 497
life table, 228
Liposcelis tricolor, 228
Opius caricivorae, 425
Opuntia cacti, 497
temperature, 228
developmental time,
cactophilic *Drosophila*, 490
host plants, 490
phenotypic plasticity, 490
viability, 490
diagnostic markers,
biological control, 749
glassy-winged sharpshooter, 749
Gonatocerus walkerjonesi, 749
natural enemies, 749
Diaprepes abbreviatus,
biological control, 413
egg parasitoid, 413
Dichroplus pratensis,
Bergmann's rule, 850
clinal variation, 850
Rensch's rule, 850
sexual size dimorphism, 850
Dipsacus,
biological control of weeds, 157
Eriophyidae, 157
Leipothrix, 157
systematics, 157
Diptera,
Acroceridae, 152
Baltic amber, 152
fossil, 152
Nemestrinoidea, 152
dispersal,
biological control, 506
Coptotermes formosanus, 506
Gloeophyllum trabeum, 506
suppression, 506
diving beetle,
larva, 787
Neotropical, 787
dual culture techniques,
Coptotermes formosanus, 703
Gloeophyllum trabeum, 703
gut cultures, 703
integument cultures, 703
Dytiscidae,
Coleoptera, 655
Laccornellus, 655
larvae, 655
phylogeny, 655

E

edge effect,
avian predation, 729
fleeing behavior, 729
habitat structure, 729
urbanization, 729
egg deposition,
cotton, 745
insect rearing, 745
soybean, 745
stink bug, 745
egg hatch,
gypsy moth, 439
inherited sterility, 439
Lymantria dispar, 439
sentinel eggs, 439
egg maturation,
bacteriocytes, 736
Bemisia aff. *gigantea*, 736
endosymbionts, 736
pharate adults, 736
egg parasitoid, 413, 640
Anagrus nigriventris, 640
biological control, 413
Diaprepes abbreviatus, 413
planthopper, 640
prairie potholes, 640
Scolochloa festucacea, 640
egg recognition,
host–parasite interaction, 532
insect–fungus interaction, 532
mimicry, 532
parasite distribution, 532
egg–pupal parasitoid,
biocontrol, 603
host location, 603
Tephritidae, 603
electronic monitoring of insect feeding,
EPG, 296
plant bug, 296
probing, 296
stylet penetration, 296
Ematurgina,
Minstrellus, 773
Neotropics, 773
phylogeny, 773
Pseudonymphidia, 773
Encyrtidae,
Georgia, 683
parasitoids, 683
Syrphophagus eliavae, 683
endophyte,
chestnut, 236
Curculio, 236
Ormyrus, 236
Torymus, 236
endosymbiont,
biological control, 275
granivore, 275
insect, 275
endosymbionts, 345, 736
bacteriocytes, 736
Baumannia, 345
Bemisia aff. *gigantea*, 736
egg maturation, 736
glassy-winged sharpshooter, 345
Homalodisca, 345
pharate adults, 736
Wolbachia, 345
engorgement,
cholecystokinin, 251
neuropeptides, 251
satiety, 251
sulfakinins, 251
Entomobryidae,
Collembola, 36
Mexico, 36
new species, 36
Willowsia, 36
EPG,
electronic monitoring of insect feeding, 296
plant bug, 296
probing, 296
stylet penetration, 296
Eriophyidae,
biological control of weeds, 157
Dipsacus, 157
Leipothrix, 157
systematics, 157

Erythroneura,
Anagrus, 41
blackberry, 41
grape, 41
leafhoppers, 41
European honey bees,
genetic markers, 64
genomic locations, 64
PCR, 64
restriction fragment length polymorphism, 64
eusociality,
positive feedback, 327
social behavior, 327
Euthamia,
gall midges, 539
monogeny, 539
natural enemies, 539
Solidago, 539
evolution,
genitalia, 100
molecular phylogeny, 100
nuclear DNA, 100
Pterostichini, 100
exotic species, 19, 474
Canary Islands, 19
geographic distribution, 474
leafhoppers, 19
Madeira, 19
Rhagoletis zephyria, 474
Sophonia orientalis, 19
Symphoricarpos spp., 474
external morphology,
harvestmen, 574
Opiliones, 574
Phalangium opilio, 574
scanning electron microscopy, 574

F

fall armyworm,
haplotype, 394
Spodoptera frugiperda, 394
fecundity,
cactus moth, 497
development, 497
life history, 497
Opuntia cacti, 497
feeding,
allogrooming, 763
social interactions, 763
subterranean termite, 763
worker behavior, 763
feeding deterrent,
Anza Borrego Desert State Park, 320
brittlebush, 320
Forelius pruinosus, 320
Linepithema humile, 320
field crop,
biological control, 876
invasive species, 876
pitfall trapping, 876
predator communities, 876
fire ant, 470, 512
biological control, 470, 512
decapitating fly, 470
geographical distribution, 512
parasitoid, 470
Pseudacteon obtusus, 512
Solenopsis gayi, 512
South America, 470
fleeing behavior,
avian predation, 729
edge effect, 729
habitat structure, 729
urbanization, 729
flight muscle,
cysteine protease inhibitor, 270
red imported fire ant queens, 270
flixweed,
Ceutorhynchus subpubescens, 178
Descurainia sophia, 178
larval taxonomy, 178
stem mining, 178
Forelius pruinosus,
Anza Borrego Desert State Park, 320
brittlebush, 320
feeding deterrent, 320
Linepithema humile, 320
fossil,
Acroceridae, 152
Baltic amber, 152
Diptera, 152
Nemestrinoidea, 152
*fts*Z,
corn rootworm, 522
mtDNA, 522
Wolbachia, 522
wsp, 522

G

gall midges,
Euthamia, 539
monogeny, 539
natural enemies, 539
Solidago, 539
gall size,
galling cynipid, 869
inquiline, 869
LMC, 869
sex ratio, 869
galling cynipid,
gall size, 869
inquiline, 869
LMC, 869
sex ratio, 869
gemmata species group,
Helotidae, 623
new species, 623
revision, 623
genetic markers,
European honey bees, 64
genomic locations, 64
PCR, 64
restriction fragment length polymorphism, 64
genitalia, 100, 403
cryptic species, 403
evolution, 100
geographic variation, 403
molecular phylogeny, 100
morphometrics, 403
mtDNA, 403
nuclear DNA, 100
Pterostichini, 100
genomic locations,
European honey bees, 64
genetic markers, 64
PCR, 64
restriction fragment length polymorphism, 64
geographic distribution,
exotic species, 474
Rhagoletis zephyria, 474
Symphoricarpos spp., 474
geographic variation,
cryptic species, 403
genitalia, 403
morphometrics, 403
mtDNA, 403
geographical distribution,
biological control, 512
fire ants, 512
Pseudacteon obtusus, 512
Solenopsis gayi, 512
Georgia,
Encyrtidae, 683
parasitoids, 683
Syrphophagus eliavae, 683
Gerridae,
Heteroptera, 139
new species, 139
Ptilomera, 139
Thailand, 139
glassy-winged sharpshooter, 345, 749
Baumannia, 345
biological control, 749
diagnostic markers, 749
endosymbionts, 345
Gonatocerus walkerjonesi, 749
Homalodisca, 345
natural enemies, 749
Wolbachia, 345
global richness,
canopy, 95
Cecidomyiidae, 95
insect galls, 95
tropical rain forest, 95

Gloeophyllum trabeum, 506, 703
biological control, 506
Coptotermes formosanus, 506, 703
dispersal, 506
dual culture techniques, 703
gut cultures, 703
integument cultures, 703
suppression, 506
Gonatocerus walkerjonesi,
biological control, 749
diagnostic markers, 749
glassy-winged sharpshooter, 749
natural enemies, 749
Gonzaga,
Berchmansus, 110
Leucochrysa, 110
new tribal assignments, 110
Vieira, 110
granivore,
biological control, 275
endosymbiont, 275
insect, 275
grape,
Anagrus, 41
blackberry, 41
Erythroneura, 41
leafhoppers, 41
grasshopper,
mate guarding, 591
polyandry, 591
protandry, 591
Sphenarium purpurascens, 591
gut cultures,
Coptotermes formosanus, 703
dual culture techniques, 703
Gloeophyllum trabeum, 703
integument cultures, 703
Guzmania,
Puerto Rico, 617
Tanytarsus bromeliad, 617
gypsy moth,
egg hatch, 439
inherited sterility, 439
Lymantria dispar, 439
sentinel eggs, 439

H

habitat structure,
avian predation, 729
edge effect, 729
fleeing behavior, 729
urbanization, 729
Halyini,
Brochymena, 673
male genitalia, 673
Parabrochymena, 673
haplotype,
fall armyworm, 394
Spodoptera frugiperda, 394
harvestmen,
external morphology, 574
Opiliones, 574
Phalangium opilio, 574
scanning electron microscopy, 574
Heleomyzidae,
host associations, 609
Neossos, 609
systematics, 609
Helotidae,
gemmata species group, 623
new species, 623
revision, 623
Hemiptera,
Alydidae, 830
Caribbean, 830
Heteroptera, 830
Neotropical, 830
hemolymph,
azadirachtin, 245
lipid metabolism, 245
Ostrinia furnacalis, 245
protein expression, 245
Hesperotyphlus,
Iberian peninsula, 444
Staphylinidae, 444
taxonomy, 444
Heteroptera, 139, 830
Alydidae, 830
Caribbean, 830
Gerridae, 139
Hemiptera, 830
Neotropical, 830
new species, 139
Ptilomera, 139
Thailand, 139
hickory,
Carya, 582
molecular systematics, 582
North America, 582
pest species, 582
Homalodisca,
Baumannia, 345
endosymbionts, 345
glassy-winged sharpshooter, 345
Wolbachia, 345
host associations,
Heleomyzidae, 609
Neossos, 609
systematics, 609
host location,
biocontrol, 603
egg–pupal parasitoid, 603
Tephritidae, 603
host plant effect,
aggregation pheromone, 83
beetle size, 83
pheromone production, 83
plant species, 83
host plants,
cactophilic *Drosophila*, 490
developmental time, 490
phenotypic plasticity, 490
viability, 490
host–parasite interaction,
egg recognition, 532
insect–fungus interaction, 532
mimicry, 532
parasite distribution, 532
hybridization,
Allonemobius, 721
Orthoptera, 721
sympatry, 721
temporal partitioning, 721
hydrolyzed yeast,
Ceratitis capitata, 900
liquid diet, 900
whole cell yeast, 900

I

Iberian peninsula,
Hesperotyphlus, 444
Staphylinidae, 444
taxonomy, 444
Idiomacromerus,
Chalcidoidea, 381
Iraella, 381
opium poppy, 381
Torymidae, 381
immature stage, 425, 914
bee nest, 914
behavior, 425
Cerioidini, 914
development, 425
Opius caricivorae, 425
sap run, 914
tree hole, 914
indirect effect,
leaf deformation, 69
Neoseiulus womersleyi, 69
Tetranychus kanzawai, 69
webs, 69
inheritance,
color polymorphism, 433
southern green stink bug, 433
X-linked trait, 433
inherited sterility,
egg hatch, 439
gypsy moth, 439
Lymantria dispar, 439
sentinel eggs, 439
inquiline,
gall size, 869
galling cynipid, 869
LMC, 869
sex ratio, 869
insect,
biological control, 275
endosymbiont, 275
granivore, 275

insect galls,
canopy, 95
Cecidomyiidae, 95
global richness, 95
tropical rain forest, 95
insect rearing,
cotton, 745
egg deposition, 745
soybean, 745
stink bug, 745
insect–fungus interaction,
egg recognition, 532
host–parasite interaction, 532
mimicry, 532
parasite distribution, 532
insect–plant interactions,
Anisoscelini, 907
Passifloraceae, 907
scanning electron microscopy, 907
stylet penetration, 907
instantaneous growth rate,
Apis mellifera, 187
resistance, 187
Russian honey bees, 187
Varroa destructor, 187
integument cultures,
Coptotermes formosanus, 703
dual culture techniques, 703
Gloeophyllum trabeum, 703
gut cultures, 703
invasive species,
biological control, 876
field crop, 876
pitfall trapping, 876
predator communities, 876
Iraella,
Chalcidoidea, 381
Idiomacromerus, 381
opium poppy, 381
Torymidae, 381
island ecology,
colonization, 711
limited mobility, 711
Ithomiinae,
Andes, 449
taxonomy, 449

J

Jiulongshan Formation,
Catiniidae, 666
new genus, 666
new species, 666
Yixian Formation, 666
juvenile hormone,
dealation, 257
nuptial flight, 257
precocene, 257

K

key for identification,
acoustic, 483
biology, 483
new species, 483
taxonomy, 483

L

Laccornellus,
Coleoptera, 655
Dytiscidae, 655
larvae, 655
phylogeny, 655
larvae, 655, 787
Coleoptera, 655
diving beetle, 787
Dytiscidae, 655
Laccornellus, 655
Neotropical, 787
phylogeny, 655
larval taxonomy,
Ceutorhynchus subpubescens, 178
Descurainia sophia, 178
flixweed, 178
stem mining, 178
leaf deformation,
indirect effect, 69
Neoseiulus womersleyi, 69
Tetranychus kanzawai, 69
webs, 69
leafhoppers, 19, 41
Anagrus, 41
blackberry, 41
Canary Islands, 19
Erythroneura, 41
exotic species, 19
grape, 41
Madeira, 19
Sophonia orientalis, 19
Leipothrix,
biological control of weeds, 157
Dipsacus, 157
Eriophyidae, 157
systematics, 157
Leucochrysa,
Berchmansus, 110
Gonzaga, 110
new tribal assignments, 110
Vieira, 110
Leucosphyrus Group,
Anopheles, 27
COI, 27
ND6, 27
phylogeny, 27
Libellulidae,
Odonata, 688
Sympetrinae, 688
Sympetrum, 688
taxonomy, 688
life history, 497, 549
cactus moth, 497
development, 497
fecundity, 497
oosorption, 549
Opuntia cacti, 497
oviposition, 549
proovigeny, 549
senescence, 549
life table,
development, 228
Liposcelis tricolor, 228
temperature, 228
limited mobility,
colonization, 711
island ecology, 711
Linepithema humile,
Anza Borrego Desert State Park, 320
brittlebush, 320
feeding deterrent, 320
Forelius pruinosus, 320
lipid metabolism,
azadirachtin, 245
hemolymph, 245
Ostrinia furnacalis, 245
protein expression, 245
Liposcelis tricolor,
development, 228
life table, 228
temperature, 228
liquid diet,
Ceratitis capitata, 900
hydrolyzed yeast, 900
whole cell yeast, 900
LMC,
gall size, 869
galling cynipid, 869
inquiline, 869
sex ratio, 869
Lycosidae,
Araneae, 1
Arctosa, 1
molecular phylogeny, 1
wolf spider, 1
Lymantria dispar,
egg hatch, 439
gypsy moth, 439
inherited sterility, 439
sentinel eggs, 439

M

Madagascar, 173, 311
mass provisioning, 311
Oligoneuriinae, 173
phylogenetics, 311
reproductive skew, 311
Rianilaneuria diminuta, 173
sociality, 311
Madeira,
Canary Islands, 19
exotic species, 19

leafhoppers, 19
Sophonia orientalis, 19
male genitalia,
Brochymena, 673
Halyini, 673
Parabrochymena, 673
mass provisioning,
Madagascar, 311
phylogenetics, 311
reproductive skew, 311
sociality, 311
mate guarding,
grasshopper, 591
polyandry, 591
protandry, 591
Sphenarium purpurascens, 591
mating,
bacteria, 418
oviposition, 418
primary reproductives, 418
ultrastructure, 418
mating behavior, 73, 333
Auchenorrhyncha, 73
chemoreception, 333
chorus, 73
contact pheromone, 333
nocturnal activity, 73
Prionus californicus, 333
vibrational communication, 73
mating success,
Danaus plexippus, 339
wing color, 339
Megachilidae,
Apidae, 810
biogeography, 810
Centriacarus, 810
Chaetodactylidae, 810
Megamelus,
Megamelus bellicus, 798
Pontederiaceae, 798
South America, 798
systematics, 798
Megamelus bellicus,
Megamelus, 798
Pontederiaceae, 798
South America, 798
systematics, 798
Mexico,
Collembola, 36
Entomobryidae, 36
new species, 36
Willowsia, 36
micronutrients,
ant, 893
Myrmica, 893
Reticulitermes, 893
termite, 893
mimicry,
egg recognition, 532
host–parasite interaction, 532
insect–fungus interaction, 532
parasite distribution, 532
Minstrellus,
Ematurgina, 773
Neotropics, 773
phylogeny, 773
Pseudonymphidia, 773
mitochondrial DNA, 289, 375
Africanized honey bee, 289
Apis mellifera, 289
cytochrome *b*, 289
mitotype, 289
phylogenetics, 375
phytophagy, 375
Schizocerella pilicornis, 375
speciation, 375
mitotype,
Africanized honey bee, 289
Apis mellifera, 289
cytochrome *b*, 289
mitochondrial DNA, 289
Mohunia,
Cicadellidae, 366
Mukariinae, 366
new genera, 366
new species, 366
molecular phylogeny, 1, 100
Araneae, 1
Arctosa, 1
evolution, 100
genitalia, 100
Lycosidae, 1
nuclear DNA, 100
Pterostichini, 100
wolf spider, 1
molecular systematics,
Carya, 582
hickory, 582
North America, 582
pest species, 582
monogeny,
Euthamia, 539
gall midges, 539
natural enemies, 539
Solidago, 539
monogyny,
oviposition, 758
polygyny, 758
Solenopsis invicta, 758
sting apparatus, 758
morphometrics,
cryptic species, 403
genitalia, 403
geographic variation, 403
mtDNA, 403
mtDNA, 403, 522
corn rootworm, 522
cryptic species, 403
ftsZ, 522
genitalia, 403
geographic variation, 403
morphometrics, 403
Wolbachia, 522
wsp, 522
Mukariinae,
Cicadellidae, 366
Mohunia, 366
new genera, 366
new species, 366
mushroom growth pattern,
competition, 222
mycophagous insect community, 222
Nitidulidae, 222
preference-performance linkage, 222
mutualisms,
ant–aphid interactions, 887
predation, 887
mycophagous insect community,
competition, 222
mushroom growth pattern, 222
Nitidulidae, 222
preference-performance linkage, 222
Myrmica,
ant, 893
micronutrients, 893
Reticulitermes, 893
termite, 893

N

natural enemies, 196, 539, 749
Aphis glycines, 196
biological control, 196, 749
diagnostic markers, 749
Euthamia, 539
gall midges, 539
glassy-winged sharpshooter, 749
Gonatocerus walkerjonesi, 749
monogeny, 539
Solidago, 539
ND6,
Anopheles, 27
COI, 27
Leucosphyrus Group, 27
phylogeny, 27
Nemestrinoidea,
Acroceridae, 152
Baltic amber, 152
Diptera, 152
fossil, 152
Neoseiulus womersleyi,
indirect effect, 69
leaf deformation, 69
Tetranychus kanzawai, 69
webs, 69
Neossos,
Heleomyzidae, 609
host associations, 609
systematics, 609
Neotropical, 787, 830
Alydidae, 830
Caribbean, 830
diving beetle, 787

Hemiptera, 830
Heteroptera, 830
larva, 787
Neotropics,
Ematurgina, 773
Minstrellus, 773
phylogeny, 773
Pseudonymphidia, 773
nest,
Apoidea, 350
bees, 350
trap-nest, 350
neuropeptides,
cholecystokinin, 251
engorgement, 251
satiety, 251
sulfakinins, 251
new genera,
Cicadellidae, 366
Mohunia, 366
Mukariinae, 366
new species, 366
new genus, 9, 666
Carnidae, 9
Catiniidae, 666
Jiulongshan Formation, 666
new species, 666
phylogeny, 9
protandrium, 9
symmetry, 9
Yixian Formation, 666
new record,
Coleotroctellus, 359
new species, 359
Taiwan, 359
Troctopsocidae, 359
new species, 36, 139, 359, 366, 390, 483, 623, 666
acoustic, 483
Aspidobracon, 390
biology, 483
Braconidae, 390
Catiniidae, 666
China, 390
Cicadellidae, 366
Coleotroctellus, 359
Collembola, 36
Entomobryidae, 36
gemmata species group, 623
Gerridae, 139
Helotidae, 623
Heteroptera, 139
Jiulongshan Formation, 666
key for identification, 483
Mexico, 36
Mohunia, 366
Mukariinae, 366
new genera, 366
new genus, 666
new record, 359
Philomacroploea, 390
Ptilomera, 139
revision, 623
Taiwan, 359
taxonomy, 483
Thailand, 139
Troctopsocidae, 359
Willowsia, 36
Yixian Formation, 666
new tribal assignments,
Berchmansus, 110
Gonzaga, 110
Leucochrysa, 110
Vieira, 110
Nitidulidae,
competition, 222
mushroom growth pattern, 222
mycophagous insect community, 222
preference-performance linkage, 222
nocturnal activity,
Auchenorrhyncha, 73
chorus, 73
mating behavior, 73
vibrational communication, 73
North America,
Carya, 582
hickory, 582
molecular systematics, 582
pest species, 582
nuclear DNA,
evolution, 100
genitalia, 100
molecular phylogeny, 100
Pterostichini, 100
nuptial flight,
dealation, 257
juvenile hormone, 257
precocene, 257

O

Odonata,
Libellulidae, 688
Sympetrinae, 688
Sympetrum, 688
taxonomy, 688
oligolecty,
Apinae, 596
bees, 596
Centridini, 596
pollination, 596
Oligoneuriinae,
Madagascar, 173
Rianilaneuria diminuta, 173
oosorption,
life history, 549
oviposition, 549
proovigeny, 549
senescence, 549
Opiliones,
external morphology, 574
harvestmen, 574
Phalangium opilio, 574
scanning electron microscopy, 574
opium poppy,
Chalcidoidea, 381
Idiomacromerus, 381
Iraella, 381
Torymidae, 381
Opius caricivorae,
behavior, 425
development, 425
immature stage, 425
Opuntia cacti,
cactus moth, 497
development, 497
fecundity, 497
life history, 497
Ormyrus,
chestnut, 236
Curculio, 236
endophyte, 236
Torymus, 236
Orthoptera, 283, 721
Allonemobius, 721
body size-related traits, 283
hybridization, 721
pericentric inversion, 283
sexual selection, 283
sympatry, 721
temporal partitioning, 721
Ostrinia furnacalis,
azadirachtin, 245
hemolymph, 245
lipid metabolism, 245
protein expression, 245
oviposition, 418, 549, 758
bacteria, 418
life history, 549
mating, 418
monogyny, 758
oosorption, 549
polygyny, 758
primary reproductives, 418
proovigeny, 549
senescence, 549
Solenopsis invicta, 758
sting apparatus, 758
ultrastructure, 418

P

Parabrochymena,
Brochymena, 673
Halyini, 673
male genitalia, 673
parasite distribution,
egg recognition, 532
host–parasite interaction, 532
insect–fungus interaction, 532
mimicry, 532
parasitoid, 470, 683
biological control, 470
decapitating fly, 470
Encyrtidae, 683

fire ant, 470
Georgia, 683
South America, 470
Syrphophagus eliavae, 683
parasitoid development,
Ceutorhynchus neglectus, 861
Ceutorhynchus obstrictus, 861
courtship behavior, 861
spatio-temporal distribution, 861
Passifloraceae,
Anisoscelini, 907
insect–plant interactions, 907
scanning electron microscopy, 907
stylet penetration, 907
PCR,
European honey bees, 64
genetic markers, 64
genomic locations, 64
restriction fragment length polymorphism, 64
pericentric inversion,
body size-related traits, 283
Orthoptera, 283
sexual selection, 283
pest species,
Carya, 582
hickory, 582
molecular systematics, 582
North America, 582
Phalangium opilio,
external morphology, 574
harvestmen, 574
Opiliones, 574
scanning electron microscopy, 574
pharate adults,
bacteriocytes, 736
Bemisia aff. *gigantea*, 736
egg maturation, 736
endosymbionts, 736
phenotypic plasticity,
cactophilic *Drosophila*, 490
developmental time, 490
host plants, 490
viability, 490
pheromone production,
aggregation pheromone, 83
beetle size, 83
host plant effect, 83
plant species, 83
Philomacroploea,
Aspidobracon, 390
Braconidae, 390
China, 390
new species, 390
phylogenetic analysis,
18S rDNA, 649
restriction fragment-length polymorphism, 649
sunflower seed weevil, 649
phylogenetics, 311, 375
Madagascar, 311
mass provisioning, 311
mitochondrial DNA, 375
phytophagy, 375
reproductive skew, 311
Schizocerella pilicornis, 375
sociality, 311
speciation, 375
phylogeny, 9, 27, 655, 773
Anopheles, 27
Carnidae, 9
COI, 27
Coleoptera, 655
Dytiscidae, 655
Ematurgina, 773
Laccornellus, 655
larvae, 655
Leucosphyrus Group, 27
Minstrellus, 773
ND6, 27
Neotropics, 773
new genus, 9
protandrium, 9
Pseudonymphidia, 773
symmetry, 9
phytophagy,
mitochondrial DNA, 375
phylogenetics, 375
Schizocerella pilicornis, 375
speciation, 375
pitfall trapping,
biological control, 876
field crop, 876
invasive species, 876
predator communities, 876
plant bug,
electronic monitoring of insect feeding, 296
EPG, 296
probing, 296
stylet penetration, 296
plant species,
aggregation pheromone, 83
beetle size, 83
host plant effect, 83
pheromone production, 83
planthopper,
Anagrus nigriventris, 640
egg parasitoid, 640
prairie potholes, 640
Scolochloa festucacea, 640
pollination,
Apinae, 596
bees, 596
Centridini, 596
oligolecty, 596
polyandry,
grasshopper, 591
mate guarding, 591
protandry, 591
Sphenarium purpurascens, 591
polygyny,
monogyny, 758
oviposition, 758
Solenopsis invicta, 758
sting apparatus, 758
Pontederiaceae,
Megamelus, 798
Megamelus bellicus, 798
South America, 798
systematics, 798
positive feedback,
eusociality, 327
social behavior, 327
prairie potholes,
Anagrus nigriventris, 640
egg parasitoid, 640
planthopper, 640
Scolochloa festucacea, 640
precocene,
dealation, 257
juvenile hormone, 257
nuptial flight, 257
predation,
ant–aphid interactions, 887
mutualisms, 887
predator communities,
biological control, 876
field crop, 876
invasive species, 876
pitfall trapping, 876
preference-performance linkage,
competition, 222
mushroom growth pattern, 222
mycophagous insect community, 222
Nitidulidae, 222
primary reproductives,
bacteria, 418
mating, 418
oviposition, 418
ultrastructure, 418
Prionus californicus,
chemoreception, 333
contact pheromone, 333
mating behavior, 333
probing,
electronic monitoring of insect feeding, 296
EPG, 296
plant bug, 296
stylet penetration, 296
proovigeny,
life history, 549
oosorption, 549
oviposition, 549
senescence, 549
protandrium,
Carnidae, 9
new genus, 9
phylogeny, 9
symmetry, 9

protandry,
grasshopper, 591
mate guarding, 591
polyandry, 591
Sphenarium purpurascens, 591
protein expression,
azadirachtin, 245
hemolymph, 245
lipid metabolism, 245
Ostrinia furnacalis, 245
Pseudacteon obtusus,
biological control, 512
fire ants, 512
geographical distribution, 512
Solenopsis gayi, 512
Pseudonymphidia,
Ematurgina, 773
Minstrellus, 773
Neotropics, 773
phylogeny, 773
Pterostichini,
evolution, 100
genitalia, 100
molecular phylogeny, 100
nuclear DNA, 100
Ptilomera,
Gerridae, 139
Heteroptera, 139
new species, 139
Thailand, 139
Puerto Rico,
Guzmania, 617
Tanytarsus bromeliad, 617
pupal color,
conditional strategy, 53
pupation, 53
stemmata, 53
pupation,
conditional strategy, 53
pupal color, 53
stemmata, 53
Pycnocephalus, 164, 839
beetles, 839
Cybocephalus, 164, 839
revision, 164
scale predator, 839
taxonomy, 164, 839

R

rearing,
adaptation, 562
artificial diet, 562
Bactrocera invadens, 562
red imported fire ant queens,
cysteine protease inhibitor, 270
flight muscle, 270
Rensch's rule,
Bergmann's rule, 850
clinal variation, 850
Dichroplus pratensis, 850
sexual size dimorphism, 850
reproductive skew,
Madagascar, 311
mass provisioning, 311
phylogenetics, 311
sociality, 311
resistance,
Apis mellifera, 187
instantaneous growth rate, 187
Russian honey bees, 187
Varroa destructor, 187
restriction fragment length polymorphism,
European honey bees, 64
genetic markers, 64
genomic locations, 64
PCR, 64
restriction fragment-length polymorphism,
18S rDNA, 649
phylogenetic analysis, 649
sunflower seed weevil, 649
Reticulitermes,
ant, 893
micronutrients, 893
Myrmica, 893
termite, 893
Reticulitermes tumiceps,
biogeography, 206
chemotaxonomy, 206
speciation, 206
subterranean termite, 206
revision, 164, 623
Cybocephalus, 164
gemmata species group, 623
Helotidae, 623
new species, 623
Pycnocephalus, 164
taxonomy, 164
Rhagoletis zephyria,
exotic species, 474
geographic distribution, 474
Symphoricarpos spp., 474
Rianilaneuria diminuta,
Madagascar, 173
Oligoneuriinae, 173
risk assessment,
5-Aza-dC, 265
Achroia grisella, 265
Apanteles galleriae, 265
toxicity, 265
Russian honey bees,
Apis mellifera, 187
instantaneous growth rate, 187
resistance, 187
Varroa destructor, 187

S

sap run,
bee nest, 914
Cerioidini, 914
immature stages, 914
tree hole, 914
satiety,
cholecystokinin, 251
engorgement, 251
neuropeptides, 251
sulfakinins, 251
scale predator,
beetles, 839
Cybocephalus, 839
Pycnocephalus, 839
taxonomy, 839
scanning electron microscopy, 574, 907
Anisoscelini, 907
external morphology, 574
harvestmen, 574
insect–plant interactions, 907
Opiliones, 574
Passifloraceae, 907
Phalangium opilio, 574
stylet penetration, 907
Schizocerella pilicornis,
mitochondrial DNA, 375
phylogenetics, 375
phytophagy, 375
speciation, 375
Scolochloa festucacea,
Anagrus nigriventris, 640
egg parasitoid, 640
planthopper, 640
prairie potholes, 640
senescence,
life history, 549
oosorption, 549
oviposition, 549
proovigeny, 549
sentinel eggs,
egg hatch, 439
gypsy moth, 439
inherited sterility, 439
Lymantria dispar, 439
sex ratio,
gall size, 869
galling cynipid, 869
inquiline, 869
LMC, 869
sexual dimorphism,
basitarsal pad, 59
bean leaf beetle, 59
corn rootworm beetle, 59
sexual selection,
body size-related traits, 283
Orthoptera, 283
pericentric inversion, 283
sexual size dimorphism,
Bergmann's rule, 850
clinal variation, 850
Dichroplus pratensis, 850
Rensch's rule, 850
social behavior,
eusociality, 327
positive feedback, 327

social interactions,
allogrooming, 763
feeding, 763
subterranean termite, 763
worker behavior, 763
sociality,
Madagascar, 311
mass provisioning, 311
phylogenetics, 311
reproductive skew, 311
Solenopsis gayi,
biological control, 512
fire ants, 512
geographical distribution, 512
Pseudacteon obtusus, 512
Solenopsis invicta,
monogyny, 758
oviposition, 758
polygyny, 758
sting apparatus, 758
Solidago,
Euthamia, 539
gall midges, 539
monogeny, 539
natural enemies, 539
Sophonia orientalis,
Canary Islands, 19
exotic species, 19
leafhoppers, 19
Madeira, 19
South America, 470, 798
biological control, 470
decapitating fly, 470
fire ant, 470
Megamelus, 798
Megamelus bellicus, 798
parasitoid, 470
Pontederiaceae, 798
systematics, 798
southern green stink bug,
color polymorphism, 433
inheritance, 433
X-linked trait, 433
soybean,
cotton, 745
egg deposition, 745
insect rearing, 745
stink bug, 745
spatio-temporal distribution,
Ceutorhynchus neglectus, 861
Ceutorhynchus obstrictus, 861
courtship behavior, 861
parasitoid development, 861
speciation, 206, 375
biogeography, 206
chemotaxonomy, 206
mitochondrial DNA, 375
phylogenetics, 375
phytophagy, 375
Reticulitermes tumiceps, 206
Schizocerella pilicornis, 375
subterranean termite, 206
Sphenarium purpurascens,
grasshopper, 591
mate guarding, 591
polyandry, 591
protandry, 591
Spodoptera frugiperda,
fall armyworm, 394
haplotype, 394
Staphylinidae,
Hesperotyphlus, 444
Iberian peninsula, 444
taxonomy, 444
stem mining,
Ceutorhynchus subpubescens, 178
Descurainia sophia, 178
flixweed, 178
larval taxonomy, 178
stemmata,
conditional strategy, 53
pupal color, 53
pupation, 53
sting apparatus,
monogyny, 758
oviposition, 758
polygyny, 758
Solenopsis invicta, 758
stink bug,
cotton, 745
egg deposition, 745
insect rearing, 745
soybean, 745
stylet penetration, 296, 907
Anisoscelini, 907
electronic monitoring of insect feeding, 296
EPG, 296
insect–plant interactions, 907
Passifloraceae, 907
plant bug, 296
probing, 296
scanning electron microscopy, 907
subterranean termite, 206, 568, 763
allogrooming, 763
biogeography, 206
chemotaxonomy, 206
feeding, 763
Reticulitermes tumiceps, 206
social interactions, 763
speciation, 206
symbiont, 568
worker behavior, 763
xylanase, 568
β-xylosidase, 568
sulfakinins,
cholecystokinin, 251
engorgement, 251
neuropeptides, 251
satiety, 251
sunflower seed weevil,
18S rDNA, 649
phylogenetic analysis, 649
restriction fragment-length polymorphism, 649
suppression,
biological control, 506
Coptotermes formosanus, 506
dispersal, 506
Gloeophyllum trabeum, 506
symbiont,
subterranean termite, 568
xylanase, 568
β-xylosidase, 568
symmetry,
Carnidae, 9
new genus, 9
phylogeny, 9
protandrium, 9
sympatry,
Allonemobius, 721
hybridization, 721
Orthoptera, 721
temporal partitioning, 721
Sympetrinae,
Libellulidae, 688
Odonata, 688
Sympetrum, 688
taxonomy, 688
Sympetrum,
Libellulidae, 688
Odonata, 688
Sympetrinae, 688
taxonomy, 688
Symphoricarpos spp.,
exotic species, 474
geographic distribution, 474
Rhagoletis zephyria, 474
Syrphophagus eliavae,
Encyrtidae, 683
Georgia, 683
parasitoids, 683
systematics, 157, 609, 798
biological control of weeds, 157
Dipsacus, 157
Eriophyidae, 157
Heleomyzidae, 609
host associations, 609
Leipothrix, 157
Megamelus, 798
Megamelus bellicus, 798
Neossos, 609
Pontederiaceae, 798
South America, 798

T

Taiwan,
Coleotroctellus, 359
new record, 359
new species, 359
Troctopsocidae, 359

Tanytarsus bromeliad,
Guzmania, 617
Puerto Rico, 617
taxonomy, 164, 444, 449, 483, 688, 839
acoustic, 483
Andes, 449
beetles, 839
biology, 483
Cybocephalus, 164, 839
Hesperotyphlus, 444
Iberian peninsula, 444
Ithomiinae, 449
key for identification, 483
Libellulidae, 688
new species, 483
Odonata, 688
Pycnocephalus, 164, 839
revision, 164
scale predator, 839
Staphylinidae, 444
Sympetrinae, 688
Sympetrum, 688
temperature,
development, 228
life table, 228
Liposcelis tricolor, 228
temporal partitioning,
Allonemobius, 721
hybridization, 721
Orthoptera, 721
sympatry, 721
Tephritidae,
biocontrol, 603
egg–pupal parasitoid, 603
host location, 603
termite,
ant, 893
micronutrients, 893
Myrmica, 893
Reticulitermes, 893
Tetranychus kanzawai,
indirect effect, 69
leaf deformation, 69
Neoseiulus womersleyi, 69
webs, 69
Thailand,
Gerridae, 139
Heteroptera, 139
new species, 139
Ptilomera, 139
Torymidae,
Chalcidoidea, 381
Idiomacromerus, 381
Iraella, 381
opium poppy, 381
Torymus,
chestnut, 236
Curculio, 236
endophyte, 236
Ormyrus, 236
toxicity,
5-Aza-dC, 265
Achroia grisella, 265
Apanteles galleriae, 265
risk assessment, 265
trap-nest,
Apoidea, 350
bees, 350
nest, 350
tree hole,
bee nest, 914
Cerioidini, 914
immature stages, 914
sap run, 914
Troctopsocidae,
Coleotroctellus, 359
new record, 359
new species, 359
Taiwan, 359
tropical rain forest,
canopy, 95
Cecidomyiidae, 95
global richness, 95
insect galls, 95

U

ultrastructure,
bacteria, 418
mating, 418
oviposition, 418
primary reproductives, 418
urbanization,
avian predation, 729
edge effect, 729
fleeing behavior, 729
habitat structure, 729

V

Varroa destructor,
Apis mellifera, 187
instantaneous growth rate, 187
resistance, 187
Russian honey bees, 187
viability,
cactophilic *Drosophila*, 490
developmental time, 490
host plants, 490
phenotypic plasticity, 490
vibrational communication,
Auchenorrhyncha, 73
chorus, 73
mating behavior, 73
nocturnal activity, 73
Vieira,
Berchmansus, 110
Gonzaga, 110
Leucochrysa, 110
new tribal assignments, 110

W

webs,
indirect effect, 69
leaf deformation, 69
Neoseiulus womersleyi, 69
Tetranychus kanzawai, 69
whole cell yeast,
Ceratitis capitata, 900
hydrolyzed yeast, 900
liquid diet, 900
Willowsia,
Collembola, 36
Entomobryidae, 36
Mexico, 36
new species, 36
wing color,
Danaus plexippus, 339
mating success, 339
Wolbachia, 345, 522
Baumannia, 345
corn rootworm, 522
endosymbionts, 345
*fts*Z, 522
glassy-winged sharpshooter, 345
Homalodisca, 345
mtDNA, 522
wsp, 522
wolf spider,
Araneae, 1
Arctosa, 1
Lycosidae, 1
molecular phylogeny, 1
worker behavior,
allogrooming, 763
feeding, 763
social interactions, 763
subterranean termite, 763
wsp,
corn rootworm, 522
*fts*Z, 522
mtDNA, 522
Wolbachia, 522

X

X-linked trait,
color polymorphism, 433
inheritance, 433
southern green stink bug, 433
β-xylosidase,
subterranean termite, 568
symbiont, 568
xylanase, 568
xylanase,
subterranean termite, 568
symbiont, 568
β-xylosidase, 568

Y

Yixian Formation,
Catiniidae, 666
Jiulongshan Formation, 666
new genus, 666
new species, 666